

EFFECTIVENESS OF BUTTERFLY PEA (*CLITORIA TERNATEA*) EXTRACT AGAINST *STREPTOCOCCUS MUTANS* BACTERIAL GROWTH IN VITRO

Siti Salmiah¹, Jesica Dwiasta Octaria Mp Nainggolan²

^{1,2} Department of Pediatric Dentistry Medan, Faculty of Dentistry, University of North Sumatra, Indonesia

Email: siti.salmiah@usu.ac.id, Jesicadwiasta2810@gmail.com

ABSTRACT

Dental caries is still a major problem in dentistry and is the most common infectious disease in children. Caries is a microbiological infectious disease that can damage the hard tissue of the teeth. Caries is caused by several factors, one of which is the microorganism Streptococcus mutans. This bacterium is the main species in dental plaque that plays a role in the etiology of caries. One way to prevent caries is cleaning plaque regularly, one of which is using mouthwash. This study aims to determine the effectiveness of butterfly pea flower extract (Clitoria ternatea) on the growth of Streptococcus mutans in vitro. This type of research is laboratory experimental research using a post test only control group design approach. The samples used were butterfly pea flower extract (Clitoria ternatea) diluted with DMSO to a concentration of 100%, 50%, 25%, 12.5%, 3.125%, 6.125% and Streptococcus mutans ATCC 25175. This research was conducted using the MIC test dilution method and KBM to see the effectiveness of the antibacterial extract against the test bacteria. Analysis of research data using the one way ANOVA test ($p < 0.05$) to see significant differences between study groups with the number of test bacteria, followed by a double comparison test with the Post Hoc Least Significance Different (LSD) method to determine the average difference between treatment groups. Antibacterial effectiveness test results showed that butterfly pea flower extract (Clitoria ternatea) was effective in inhibiting Streptococcus mutans bacteria ($p < 0.05$), at a concentration of 25% was the minimum inhibitory level (MIC) and 50% concentration was the minimum killing rate (MBC). The conclusion of this study is that the extract of butterfly pea flower (Clitoria ternatea) is effective in inhibiting and killing Streptococcus mutans bacteria.

KEYWORDS

Dental Plaque, Streptococcus Mutans Butterfly Pea Flower, MIC, KBM.



This work is licensed under a Creative Commons Attribution-ShareAlike 4.0 International

How to cite: Siti Salmiah, Jesica Dwiasta Octaria Mp Nainggolan (2024). Effectiveness of Butterfly Pea (Clitoria Ternatea) Extract Against Streptococcus Mutans Bacterial Growth In Vitro. *Journal Eduvest*. 4 (10): 8843-8850
E-ISSN: 2775-3727
Published by: <https://greenpublisher.id/>

INTRODUCTION

Dental and oral health problems are very important in health development, especially in elementary school-age children (Sherlyta et al., 2017). Dental caries is still a major problem in dentistry and is the most common infectious disease in children. Caries is a microbiological infectious disease of the teeth that can damage the hard tissue of the teeth. The presence of a cavity in the tooth is a sign of a bacterial infection. The results of the 2018 Basic Health Research show that dental and oral problems have reached 57.6% in Indonesia. Bagramian et al in 2009 said that nearly 90% of school-age children worldwide suffer from dental caries. Meanwhile, according to the 2013 Centers of Control Disease Prevention (CDC), dental caries is a chronic disease that often occurs in children aged 6-11 years (25%) and adolescents aged 12-19 years (59%). This problem also occurs in Indonesia. In 2014, the Indonesian Ministry of Health said that 89% of children under the age of 12 suffer from dental caries. Caries is stated as a multifactorial disease, namely the presence of several main factors, namely host factors, microorganisms, substrate and time. One of them is the role of *Streptococcus mutans* which has acidogenic (acid producing) and aciduric (acid resistant) properties (Gayatri, 2017).

Streptococcus mutans is the main species in dental plaque which plays an important role in the etiology of caries. *Streptococcus mutans* has glucosyltransferase (GTF) and fructosyltransferase (FTF) enzymes which convert sucrose into glucans and fructans which help the attachment of *Streptococcus mutans* bacteria and the formation of plaque on the tooth surface. It is very important to prevent caries from an early age, one way to prevent it is by cleaning plaque regularly. Plaque cleaning can be done mechanically and chemically. Mechanically it can be done by brushing your teeth and flossing while chemically you can use toothpaste and mouthwash (Pujoraharjo & Herdiyati, 2018).

According to Fajrian, et al in 2014, the use of 0.2% chlorhexidine once a day can reduce *Streptococcus mutans* bacteria by 30-50%. However, there is a drawback to chlorhexidine, as it is a chemical that can give off an unpleasant taste and cause staining of the teeth. The return of attention to natural materials (back to nature) has long been considered a useful thing because it has been believed to prevent various kinds of diseases. Many studies have been carried out by utilizing natural materials with the aim of producing medicines in an effort to support dental health service programs, especially preventing caries (Carranza & Newman, 2018).

One of the plants that has the potential as an herbal ingredient is butterfly pea flower (*Clitoria ternatea*). This flower is increasingly popular in Indonesia as a flower that provides many health benefits. Butterfly pea flowers extracted using various solvents showed broad antimicrobial activity including gram-positive bacteria, gram negative bacteria and fungi (Marpaung, 2020). Butterfly pea flower is an herbal plant in Indonesia which provides many health benefits, where the ingredients contained in this flower contain phytochemical compounds including flavonoids and their derivatives flavonol glycosides, kaempferol glycosides,

quercetin glycosides, myricetin glycosides and anthocyanins. This compound is present in all parts of the plant including leaves, roots, wood, flowers, fruit and seeds, which are identified as antibacterial (Angriani, 2019).

Research conducted by Riyanto EF, et al in 2019 said that 70% ethanol extract of butterfly pea flowers can inhibit the growth of *Pseudomonas aeruginosa* bacteria with a concentration of 10% and *Bacillus cereus* with a concentration of 30% (Riyanto & Suhartati, 2019). Previous research was also conducted by Widyarman AS, et al in 2018 regarding the effectiveness of butterfly pea flower extract (*Clitoria ternatea*) in vitro to inhibit *Porphyromonas gingivalis* bacteria. The results of the study by Widyarman et al said that the concentration of butterfly pea flower extract that can inhibit *Porphyromonas gingivalis* bacteria is at a concentration of 50% (Widyarman et al., 2018).

Based on the background of the problems that have been described above, research on the effectiveness of butterfly pea flower extract (*Clitoria ternatea*) against *Streptococcus mutans* bacteria is still limited, so researchers are interested in researching butterfly pea flower extract (*Clitoria ternatea*) at concentrations of 100%, 50%, 25%, 12.5%, 6.25% and 3.125% which were extracted with 70% alcohol using a positive control of chlorhexidine and a negative control of Dimethylsulfoxide (DMSO) in inhibiting the growth of *Streptococcus mutans* so that later butterfly pea flowers can be used as an alternative to traditional caries prevention (Ifitri & Eriyati, 2019). This study aims to determine the antibacterial effectiveness of butterfly pea flower extract (*Clitoria ternatea*) with each concentration of 100%, 50%, 25%, 12.5%, 6.25%, 3.125% and chlorhexidine on the growth of *Streptococcus mutans* bacteria.

RESEARCH METHOD

Type of research is laboratory experimental research using a post test only control group design approach. The samples used were butterfly pea flower extract (*Clitoria ternatea*) diluted with DMSO to a concentration of 100%, 50%, 25%, 12.5%, 3.125%, 6.125% and *Streptococcus mutans* ATCC 25175. This research was conducted using the MIC test dilution method and KBM to see the effectiveness of the antibacterial extract against the test bacteria. Analysis of research data using the one way ANOVA test ($p < 0.05$) to see significant differences between study groups with the number of test bacteria, followed by a double comparison test with the Post Hoc Least Significance Different (LSD) method to determine the average difference between treatment groups (Suryani, 2019).

RESULT AND DISCUSSION

a post test only control group design research conducted in two laboratories, namely the Traditional Medicine Laboratory, USU Faculty of Pharmacy, to extract butterfly pea flower (*Clitoria ternatea*) and then test the effectiveness of the extraction results on the growth of *Streptococcus mutans* bacteria in vitro at the USU Hospital Microbiology Laboratory. This study used six treatment groups

consisting of six different concentrations of butterfly pea flower extract (*Clitoria ternatea*) namely 100%, 50%, 25%, 12.5%, 6.25% and 3.125%. Chlorhexidine 0.2% was used as a positive control and DMSO as a negative control. In this study, the dilution method was used to determine the Minimum Inhibitory Concentration (MIC) and Minimum Inhibitory Concentration (MBC) with 4 repetitions in each test tube (Yu et al., 2017).

The test tube which already contains Nutrient Broth media is added to the butterfly pea flower extract in various concentrations and then *Streptococcus mutans* suspension with 0.5 McFarland turbidity is added. The tubes were vortexed and incubated in an incubator for 24 hours at 37 °C After that, observations were made for the presence or absence of sediment on all the bottom of the tube. The tube that looks clear (bright purple) and has no precipitate with the smallest concentration of butterfly pea extract is the temporary MIC value of the butterfly pea extract on the growth of *Streptococcus mutans* bacteria.

Observation of the dilution tube has been completed, so to ensure that the tube has bacterial growth or not, it is continued to the subculture stage on TYCSB media and incubated in an incubator for 24 hours at 37°C, after 24 hours it will be observed whether or not bacterial colonies are growing on TYCSB media. The petri dish where there was no bacterial growth with the smallest concentration of butterfly pea extract was the MBC value of the butterfly pea extract on the growth of *Streptococcus mutans* bacteria (Kusuma et al., 2019).

Determination of Minimum Inhibitory Content (MIC) and Minimum Killing Content (KBM) of Butterfly Pea Flower Extract (*Clitoria ternatea*) on the Growth of *Streptococcus mutans* Bacteria. In the dilution method, the MIC and MBC values were observed after the incubation period by observing the turbidity level in each concentration tube, but the turbidity level of the butterfly pea extract could not be determined because the concentration of the butterfly pea extract was too concentrated and dark, so it could only be seen by categorizing the color purple. dark or light purple (Kolliyavar et al., 2016). Observations were made on all repetitions of each concentration of butterfly pea extract at the same time.

Based on Figure 1, it can be concluded that the Minimum Inhibitory Content (MIC) value is 25%, which at this concentration has a bright purple color, while the Minimum Inhibitory Concentration (KBM) value is found at a concentration of 50% which has a dark purple color. Furthermore, to prove the MIC and MBC values with certainty, a subculture stage was carried out in each dilution tube using TYCSB (Tryptone-yeast-cysteinesucrose-bacitracin) media. Subculture on TYCSB media was carried out to obtain MIC and MBC values of *Streptococcus mutans* ATCC 25175. Calculation of the number of bacterial colonies on a petri dish was carried out using the TPC (Total Plate Count) method by means of visual observation and counting (Purba, 2020).

In table 1, it can be seen that the results of dilution tube culture at the lowest concentration that can inhibit bacterial growth and show a bacteriostatic effect is a concentration of 25% with an average value of 37 CFU/ml, then this concentration is determined as the MIC value. The concentration of 50% is the lowest concentration where there is no bacterial growth and shows a bacteriocidal effect is

a concentration of 50% with an average value of the number of bacterial colonies of 0 CFU/ml, then this concentration is determined as the MBC value.

Antibacterial Effectiveness Test of Each Concentration of Butterfly Pea Flower (*Clitoria ternatea*) Extract on the Growth of *Streptococcus mutans* Bacteria
Antibacterial effectiveness of each concentration of butterfly pea flower extract (*Clitoria ternatea*) against *Streptococcus mutans* was analyzed using the One Way ANOVA test. Before the One Way ANOVA test was carried out the data normality test was first carried out using the Shapiro-Wilk test to find out whether the data was normally distributed or not. If the data is normally distributed, then the test is continued by using the One Way ANOVA and LSD (Least Significant Difference) tests.

The results of the Shapiro-Wilk normality test obtained data at a concentration of 25% with a value of $p = 0.319$, a concentration of 12.5% with a value of $p = 0.934$, a concentration of 6.25% with a value of $p = 0.636$, and a concentration of 3.125% with a value of $p = 0.448$ ($p > 0.05$). This shows that the data at each concentration is normally distributed because all p values are > 0.05 , so that the One Way ANOVA test can be carried out. In table 3, a significance of $p = 0.000$ ($p < 0.05$) is obtained, which means that there is antibacterial effectiveness of butterfly pea flower extract (*Clitoria ternatea*) on the growth of *Streptococcus mutans* bacteria. After the One Way ANOVA test was carried out, the LSD (Least Significant Difference) test was then performed to find out which treatment group pairs were significant. The LSD (Least Significant Difference) test results showed that there was a statistically significant difference between each concentration group.

Discussion

Streptococcus mutans bacteria was chosen as the research sample because *Streptococcus mutans* is one of the most caries-causing microbial pathogens found in plaque, and in children with high levels of caries, there will be an increase in the number of *Streptococcus mutans* colonies in the oral cavity.² Researchers used butterfly pea extract in this study because the butterfly pea flower has the ability to inhibit the growth of several types of bacteria.⁸ This study used butterfly pea flower (*Clitoria ternatea*) extract with 70% ethanol at concentrations of 100%, 50%, 25%, 12.5%, 6.25%, and 3.125% and used pure *Streptococcus mutans* isolates. The isolate was obtained from the American Type Culture Collection (ATCC) numbered 25175.

the one way ANOVA statistical test namely the effectiveness test of butterfly pea flower extract (*Clitoria ternatea*) with concentrations of 100%, 50%, 25%, 12.5%, 6.25%, and 3.125% on the growth of *Streptococcus mutans* ATCC 25175 in an vitro has a significance value of $p = 0.000$ ($p < 0.05$). This value indicates the presence of inhibition of butterfly pea flower extract which has an antibacterial effect on the growth of *Streptococcus mutans* ATCC 25715 in vitro To see significant differences between treatment groups, a posthoc Least Significant Difference (LSD) statistical test was performed.

The results of this study prove that butterfly pea flower extract (*Clitoria ternatea*) has antibacterial activity. This effect can occur because the butterfly pea flower (*Clitoria ternatea*) has a compound that has the potential as an antibacterial,

namely flavonoids. According to research by Sankari et al in 2014, the tested flavonoids have antibacterial activity, especially against *Streptococcus mutans* 31 Flavonoids cause damage to the permeability of bacterial cell walls, microsomes, and lysosomes as a result of interactions between flavonoids and bacterial DNA. 40 Flavonoids contained in the butterfly pea flower are 20.07 ± 0.55 mmol/mg. 27 The purple color of the butterfly pea flower indicates the presence of anthocyanin compounds, so that in some countries the butterfly pea flower is used as a natural food coloring. Nigam and Shivastava in Riyanto EF, et al in 2019 stated that phytochemical compounds such as alkaloids, flavonoids and others can be useful as antibacterials against the growth of many microorganisms.10 Phytochemical compounds that are antibacterial are effective against the growth of gram positive and gram-negative bacteria. This can be seen from previous studies regarding the effectiveness of butterfly pea flower extract as an antibacterial.

This research is in line with the research of Pratap et al in 2012 which stated that butterfly pea flower extract (*Clitoria ternatea*) used Distilled water as a solvent against *Streptococcus mutans*, *Lactobacillus casei*, *Staphylococcus aureus* the most effective concentration was 50% where the inhibition zone formed from *Streptococcus mutans* 7 mm, *Lactobacillus casei* 8 mm, *Staphylococcus aureus* 10 mm, so it was found that *Staphylococcus aureus* had the largest inhibition zone.

Another previous study was also conducted by Widyarman et al in 2018 stating that the extract of butterfly pea flower juice (*Clitoria ternatea*) was effective in inhibiting the growth of *Porphyromonas gingivalis* bacteria at a concentration of 50%. 11 Another study was also conducted by Riyanto EF et al in 2019 said that 70% ethanol extract of butterfly pea flowers can inhibit the growth of *Pseudomonas aeruginosa* bacteria with a minimum inhibitory concentration of 10%, and can inhibit the growth of *Bacillus cereus* bacteria with a minimum inhibitory concentration of 30%.10

Differences in the results of the antibacterial effect from previous studies which had natural ingredients against *Streptococcus mutans* bacteria could have been caused by differences in methods, differences in materials used, differences in solvents used, differences in extraction methods, and differences in the types of active compounds of each of these natural ingredients.

The results of this study also found that the MIC value of the butterfly pea flower extract test was not able to exceed the value of the positive control test (0.2% chlorhexidine) because 0.2% chlorhexidine contained a stable bisbiguanide compound. 42 In contrast to the flavonoid compounds found in butterfly pea flower (*Clitoria ternatea*), which are unstable. Unstable flavonoid compounds can occur because they can be influenced by the solvent used or the temperature during extraction 42 Plaque inhibition can occur due to the presence of flavonoid compounds in butterfly pea flowers which can inactivate the glucosyltransferase enzyme which plays an important role in plaque formation.

Caries will easily occur if plaque formation is not inhibited because *Streptococcus mutans* also produces extracellular polysaccharides which can facilitate plaque attachment to the tooth surface and bacteria will produce acid which can affect demineralization.45

This study also proves that the hypothesis proposed by the researcher is acceptable, namely that the butterfly pea flower extract with 70% ethanol solvent can inhibit plaque formation and inhibit the *Streptococcus mutans* bacteria in the oral cavity because it contains flavonoids, alkaloids and anthocyanins, so research on butterfly pea extract (*Clitoria ternatea*) Later it can be used as a reference for research on butterfly pea flower (*Clitoria ternatea*) with mouthwash preparations or topical preparations for young children because it can inhibit *Streptococcus mutans* bacteria in the oral cavity at a concentration of 25% and kill *Streptococcus mutans* bacteria at a concentration of 50%.

CONCLUSION

The conclusion of this study is that telang flower extract (*Clitoria ternatea*) is effective in inhibiting and killing *Streptococcus mutans* bacteria that play a role in the formation of dental caries. At a concentration of 25%, telang flower extract showed the ability to inhibit bacterial growth (*Minimum Inhibitory Concentration* or MIC), while at a concentration of 50%, the extract was able to kill bacteria (*Minimum Bactericidal Concentration* or MBC). Flavonoids, alkaloids, and anthocyanins contained in telang flowers are believed to be the main components that contribute to this antibacterial activity. This study shows that bayang flowers have potential as a natural alternative for caries prevention, with effectiveness that approaches, although does not exceed, the positive control of chlorhexidine 0.2%.

REFERENCES

- Angriani, L. (2019). Potensi Ekstrak Bunga Telang (*Clitoria Ternatea*) Sebagai Pewarna Alami Lokal Pada Berbagai Industri Pangan:(The Potential of Extract Butterfly Pea Flower (*Clitoria ternatea* L.) as a Local Natural Dye for Various Food Industry). *Canrea Journal: Food Technology, Nutritions, and Culinary Journal*, 32–37.
- Carranza, F. A., & Newman, M. (2018). *Carranza's Clinical Periodontology*.
- Gayatri, R. W. (2017). Hubungan Tingkat Pengetahuan dengan Perilaku pemeliharaan Kesehatan Gigi anak SDN KAUMAN 2 MALANG. *Journal of Health Education*, 2(2), 194–203.
- Ifitri, I., & Eriyati, E. (2019). PENGARUH BERKUMUR AIR REBUSAN DAUN JAMBU BIJI TERHADAP INDEKS PLAK PADA MURID SDN 10 KUBU AMPEK ANGKEK KABUPATEN AGAM. *Ensiklopedia of Journal*, 1(4).
- Kolliyavar, B., Shettar, L., & Thakur, S. (2016). Chlorhexidine: the gold standard mouth wash. *J Pharm Biomed Sci*, 6(2), 106–109.
- Kusuma, Y., Pinatih, K. J. P., & Hendrayana, M. A. (2019). Efek sinergis kombinasi chlorhexidine dan alkohol terhadap daya hambat pertumbuhan *Staphylococcus aureus*. *E-Jurnal Medika*, 8(3), 139–146.
- Marpaung, A. M. (2020). Tinjauan manfaat bunga telang (*clitoria ternatea* l.) bagi kesehatan manusia. *Journal of Functional Food and Nutraceutical*, 63–85.
- Pujoraharjo, P., & Herdiyati, Y. (2018). Efektivitas antibakteri tanaman herbal terhadap *streptococcus mutans* pada karies anak. *Indonesian Journal of Paediatric Dentistry*, 1(1), 51–56.

- Purba, E. C. (2020). Kembang telang (*Clitoria ternatea* L.): pemanfaatan dan bioaktivitas. *Jurnal EduMatSains*, 4(2), 111–124.
- Riyanto, E. F., & Suhartati, R. (2019). Daya hambat ekstrak etanol bunga telang (*Clitoria ternatea* L) terhadap bakteri perusak pangan. *Jurnal Kesehatan Bakti Tunas Husada: Jurnal Ilmu-Ilmu Keperawatan, Analisis Kesehatan Dan Farmasi*, 19(2), 218–225.
- Sherlyta, M., Wardani, R., & Susilawati, S. (2017). Tingkat kebersihan gigi dan mulut siswa Sekolah Dasar Negeri di desa tertinggal Kabupaten Bandung Oral hygiene level of underdeveloped village State Elementary School students in Bandung Regency. *Jurnal Kedokteran Gigi Universitas Padjadjaran*, 29(1), 69–76.
- Suryani, N. (2019). Obat kumur herbal yang mengandung ekstrak etil asetat kulit batang bintaro (*Cerbera odollam* Gaertn) sebagai antibakteri *Streptococcus mutans* penyebab plak gigi. *Farmaka*, 17(2), 48–56.
- Widyarman, A. S., Sumadi, S., & Agustin, T. P. (2018). Antibiofilm effect of *Clitoria ternatea* flower juice on *Porphyromonas gingivalis* in vitro. *Journal of Indonesian Dental Association*, 1(1).
- Yu, O. Y., Zhao, I. S., Mei, M. L., Lo, E. C.-M., & Chu, C.-H. (2017). Dental biofilm and laboratory microbial culture models for cariology research. *Dentistry Journal*, 5(2), 21.